



YAYASAN NURUL JADID PAITON
**LEMBAGA PENERBITAN, PENELITIAN, &
PENGABDIAN KEPADA MASYARAKAT**
UNIVERSITAS NURUL JADID
PROBOLINGGO JAWA TIMUR

PP. Nurul Jadid
Karanganyar Paiton
Probolinggo 67291
☎ 0888-3077-077
e: lp3m@unuja.ac.id
w: <https://lp3m.unuja.ac.id>

SURAT KETERANGAN

Nomor : NJ-To6/0351/A.03/LP3M/02.2021

Lembaga Penerbitan, Penelitian, dan Pengabdian Masyarakat (LP3M) Universitas Nurul Jadid Probolinggo menerangkan bahwa artikel/karya tulis dengan identitas berikut ini:

Judul : *Terwilliger Algebras Of Some Group Association Schemes*
Penulis : NUR HAMID, M.Si, Ph.D..
Identitas : Mathematical Journal of Okayama University, Vol. LXI, No.1,
Januari 2019, E-ISSN : 0030-1566, Okayama University
No. Pemeriksaan : 012085

Telah selesai dilakukan *similarity check* dengan menggunakan perangkat lunak **Turnitin** pada 17 Januari 2021 dengan hasil sebagai berikut: Tingkat kesamaan diseluruh artikel (*Similarity Index*) adalah 23 % dengan publikasi yang telah diterbitkan oleh penulis pada Mathematical Journal of Okayama University (<http://www.math.okayama-u.ac.jp/mjou/>)

Demikian Surat Keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Probolinggo, 02 Februari 2021

Kepala LP3M,



ACHMAD FAWAID, M.A., M.A.

NIDN. 2123098702

TERWILLIGER ALGEBRAS OF SOME GROUP ASSOCIATION SCHEMES

Submission date: 17-Jan-2021 04:52PM (UTC-0800)

Submission ID: 1489178517

File name: artikel1_Terwilliger_okk.pdf (357.48K)

Word count: 2310

Character count: 9413

TERWILLIGER ALGEBRAS OF SOME GROUP ASSOCIATION SCHEMES

Nur HAMID AND MANABU OURA

ABSTRACT. The Terwilliger algebra plays an important role in the theory of association schemes. The present paper gives the explicit structures of the Terwilliger algebras of the group association schemes of the finite groups $PSL(2, 7)$, A_6 , and S_6 .

1. INTRODUCTION

Association schemes enable us to study combinatorial problems in a unified way. We refer to [2, 6] for the foundations of association schemes. In a series of papers [10, 11, 12], Terwilliger introduced a new method, the so-called Terwilliger algebra, to investigate the commutative association schemes. Since then there have been many investigations on Terwilliger algebras (cf. [8, 7]). It is very important to know the explicit structure of the Terwilliger algebra. The cases of the group association schemes of S_5 and A_5 were studied in [1] along the line of the work [3]. In the present paper we determine the structures of the Terwilliger algebras of the group association schemes of the finite groups $PSL(2, 7)$, A_6 , and S_6 .

The computations were done with Magma [5] and SageMath [9].

2. PRELIMINARIES

We begin with the definition of a group association scheme.

Definition 1. Let G be a finite group and $C_0 = \{e\}, C_1, \dots, C_d$ the conjugacy classes of G , where e is the identity of G . Define the relations $R_i (i = 0, 1, \dots, d)$ on G by

$$(x, y) \in R_i \iff yx^{-1} \in C_i.$$

Then $X(G) = (G, \{R_i\}_{i=0}^d)$ forms a commutative association scheme of class d called the group association scheme of G .

We associate the matrix A_i of the relation R_i as

$$(A_i)_{x,y} := \begin{cases} 1 & \text{if } (x, y) \in R_i, \\ 0 & \text{otherwise.} \end{cases}$$

Mathematics Subject Classification. Primary 35C07; Secondary 35K57.

Key words and phrases. Terwilliger algebra, group association scheme.

Then we have

$$\sum_{i=0}^d A_i = K_{ij}$$

and A_0, \dots, A_d generate the so-called Bose-Mesner algebra A . The intersection numbers p^k_{ij} of the group association scheme $X(G)$ are given

by

$$|\{(x, y) \in C_i \times C_j \mid xy = z, z \in C_k\}|.$$

The algebra A has a second basis E_0, \dots, E_d of primitive idempotents, and

$$E_i \circ E_j = \frac{1}{|G|} q^k_{ij} E_k,$$

where \circ denotes Hadamard (entry-wise) multiplication. For each $i = 0, \dots, d$, let E_i^* and A_i^* be the diagonal matrices of size $|G|$ which are defined as follows.

$$(E_i^*)_{x,x} := \begin{cases} 1, & \text{if } x \in C_i \\ 0, & \text{if } x \notin C_i \end{cases} \quad (x \in G),$$

$$(A_i^*)_{x,x} := |G|(E_i)_{e,x} \quad (x \in G).$$

Then E_0^*, \dots, E_d^* form a basis for the dual Bose-Mesner algebra A^* . The

intersection numbers provide information for our structural results to follow. We refer to the following relations [10].

$$\begin{aligned} E_i^* A_j E_k^* &= 0 \quad \text{if } ij = 0 \quad (0 \leq i, j, k \leq d), \\ \Leftrightarrow p^k_{ij} E_i A_j E_k &= 0 \quad (0 \leq i, j, k \leq d). \\ = 0 &\Leftrightarrow q^k_{ij} \end{aligned}$$

We need to fix the ordering of the conjugacy classes. The following table gives the representatives and the orders of conjugacy classes.

(1) $PSL(2, 7)$

	C_0	C_1	C_2
rep.	(1)	(357)(468)	(2354786)
$ C_i $	1	56	24
	C_3	C_4	C_5
rep.	(2465837)	(12)(34)(58)(67)	(1235)(4876)
$ C_i $	24	21	42

(2) A_6

	C_0	C_1	C_2	C_3	C_4
rep.	(1)	(12)(34)	(123)	(123)(456)	(1234)(56)
$ C_i $	1	45	40	40	90
	C_5	C_6			
rep.	(12345)	(12346)			
$ C_i $	72	72			

(3) S_6

	C_0	C_1	C_2	C_3
rep.	(1)	(12)	(12)(34)	(12)(34)(56)
$ C_i $	1	15	45	15
	C_4	C_5	C_6	C_7
rep.	(123)	(123)(45)	(123)(456)	(1234)
	40	120	40	90
	C_8	C_9	C_{10}	
	(1234)(56)	(12345)	(123456)	
	90	144	120	

Finally we give the definition of the Terwilliger algebra of the group association scheme. We shall denote by k the ring of k matrices over the complex number \mathbb{C} .

Definition 2. Let G be a finite group. The Terwilliger algebra $T(G)$ of the group association scheme $X(G)$ is a sub-algebra of $M_n(k)$ generated by A and A^* .

Since $T(G)$ is closed under the conjugate-transpose, $T(G)$ is semi-simple. In the next section, we investigate the Terwilliger algebras of the group association schemes of $PSL(2, 7)$, A_6 and S_6 .

3. RESULTS

In [1], Balmaceda and Oura gave the structures of the Terwilliger algebra of the group association schemes of S_5 and A_5 . Following their method, we determine the Terwilliger algebras for the cases $PSL(2, 7)$, A_6 , and S_6 .

Theorem 3.1. The dimensions of $T(PSL(2, 7))$, A_6 and $T(S_6)$ are given as follows.

$$\begin{aligned} \dim T(PSL(2, 7)) &= 165, \dim T(A_6) \\ &= 336, \dim T(S_6) = 758. \end{aligned}$$

Proof. We compute a set of linearly independent elements among $E_i^* A_j E_k^*$ and $E_i^* A_j E_k^* \cdot E_{k_1}^* A_l E_{m_1}^* = E_i^* A_j E_{k_1}^* A_l E_{m_1}^*$. By direct calculation we can see that any form $E_i^* A_{i_1} E_{j_1}^* \cdot E_{j_2}^* A_{i_2} E_{k_1}^* \cdot E_{k_2}^* A_{i_3} E_{l_1}^*$ linearly depends on the $E_i^* A_{i_4} E_{l_1}^*$'s and the $E_i^* A_{i_5} E_{k_1}^* \cdot E_{k_1}^* A_{i_6} E_{l_1}^*$'s. Therefore the products of more than two elements of the form $E_i^* A_j E_k^*$ give no new elements of a basis. \square

¹This answers a question raised by Prof. Terwilliger. Indeed our original argument had a gap. He informed us the reference [4]. Our result $\dim T(S_6) = 758$ is violated to Conjecture 3.5.

$$\dim Z(T(S_6)) = 14.$$

Proof. The result is obtained by determining a basis for the center. We solve a linear equation system $\{x_i y = y x_i\}$ ranging over all the basis of $T(G)$ and $c_j \overline{b_j}$, where b_j are the basis elements of G and c_j is any scalar. \square

Let $\{e_i : 1 \leq i \leq s\}$ be a basis of $Z(T(G))$. Then we have $e_i e_j = t^k e_k$ and put $B_i := (t^k)$ for $1 \leq i \leq s$. Since these matrices mutually commute, they are simultaneously diagonalizable. We shall denote by $v_1(i), \dots, v_s(i)$ the diagonal entries of the diagonalized matrix of B_i and define the matrix M by $M_{ij} := v_i(j)$. Then we get the primitive central idempotents $\varepsilon_1, \dots, \varepsilon_s$ by

$$(\varepsilon_1, \dots, \varepsilon_s) = (e_1, \dots, e_s) M^{-1}.$$

Theorem 3.3. *The degrees of the irreducible complex representations afforded by every idempotent are given below.*

$T(PSL(2, 7))$	ε_i	ε_1	ε_2	ε_3	ε_4	ε_5	ε_6	ε_7	
deg	ε_i	1	2	3	3	5	6	9	
$T(A_6)$	ε_i	ε_1	ε_2	ε_3	ε_4	ε_5	ε_6	ε_7	ε_8
deg	ε_i	1	3	3	4	4	6	6	7
	ε_i	ε_9	ε_{10}						
	ε_i	8	10						
	deg								
$T(S_6)$	ε_i	ε_1	ε_2	ε_3	ε_4	ε_5	ε_6	ε_7	ε_8
deg	ε_i	1	1	1	3	3	4	6	7
		ε_9	ε_{10}	ε_{11}	ε_{12}	ε_{13}	ε_{14}		
		8	8	9	11	15			

Proof. This is because that $T(G)\varepsilon_i \cong M_{d_i}$ and that $d_i = \dim T(G)\varepsilon_i$ equals the number of linearly independent elements in the set $\{x_j \varepsilon_i\}$, where x_j are the basis elements of T . \square

Theorems 3.1 and 3.3 are combined as $165 = 1^2 + 2^2 + 3^2 + 3^2 + 5^2 + 6^2 + 9^2$,

$$336 = 1^2 + 3^2 + 3^2 + 4^2 + 4^2 + 6^2 + 6^2 + 7^2 + 8^2 + 10^2,$$

$$758 = 1^2 + 1^2 + 1^2 + 3^2 + 3^2 + 4^2 + 6^2 + 7^2 + 8^2 + 8^2 + 9^2 + 11^2 + 15^2.$$

The degrees of irreducible complex representations afforded by every primitive central idempotents enable us to get the following structure theorem.

Corollary 3.4. *We have that*

$$T(PSL(2, 7)) \cong M_1 \oplus M_1 \oplus M_3 \oplus M_3 \oplus M_5 \oplus M_6 \oplus M_9,$$

$$\begin{aligned}
 T(A_5) &\cong M_1 \oplus M_2 \oplus M_3 \oplus M_4 \oplus M_5 \oplus M_6 \oplus M_7 \\
 &\quad \oplus M_8 \\
 &\quad \oplus M_{10}, \\
 T(S_5) &\cong M_1 \oplus M_2 \oplus M_3 \oplus M_4 \oplus M_5 \oplus M_6 \oplus M_7 \\
 &\quad \oplus M_8 \\
 &\quad \oplus M_9 \oplus M_{10} \oplus M_{11} \oplus M_{15}.
 \end{aligned}$$

Acknowledgment. The second named author was supported by JSPS KAKENHI Grant Number JP25400014. The authors would like to thank Prof. Terwilliger for helpful discussions.

REFERENCES

- [1] J. M. P. Balmaceda, M. Oura, *The Terwilliger algebras of the group association schemes of S_5 and A_5* , Kyushu J. Math., **48**, no. 3 (1994), 221 - 231.
- [2] E. Bannai, T. Ito, *Algebraic Combinatorics I: Association Schemes*, Benjamin/Cummings, California, 1984.
- [3] E. Bannai, A. Munemasa, *The Terwilliger algebras of group association schemes*, Kyushu J. Math., **49**, no. 1 (1995), 93 - 102.
- [4] M. M. Bogaerts, P. Dukes, *Semidefinite programming for permutation codes*, Discrete Mathematics, **326**, (2014), 34 - 43.
- [5] W. Bosma, J. Cannon, C. Playoust, *The Magma algebra system I: The user language*, J. Symbolic Comput., **24**, no. 3-4 (1997), 235 - 265.
- [6] A. E. Brouwer, C. M. Cohen, A. Neumaier, *Distance-regular graphs*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 18. Springer-Verlag, Berlin, 1987.
- [7] T. Ito, *Tridiagonal pairs and q -Onsager algebras (Japanese)*, SIAM J. Math. Anal., **Sūgaku** **65**, no. 1 (2013), 69 - 92.
- [8] W. Martin, H. Tanaka, *Commutative association schemes*, European J. Combin., **30**, no. 6 (2009), 1497 - 1525.
- [9] The Sage Developers, *Sagemath, the Sage Mathematics Software System (Version 6.4.1)*, <http://www.sagemath.org>, 2014.
- [10] P. Terwilliger, *The subconstituent algebra of an association scheme I*, J. Algebraic Combin., **1**, no. 4 (1992), 363 - 388.
- [11] P. Terwilliger, *The subconstituent algebra of an association scheme II*, J. Algebraic Combin., **2**, no. 1 (1993), 73 - 103.
- [12] P. Terwilliger, *The subconstituent algebra of an association scheme III*, J. Algebraic Combin., **2**, no. 2 (1992), 177 - 210.

Nur HAMID
 FACULTY of MATHEMATICS AND Physics, KANAZAWA
 UNIVERSITY AND
 NURUL JADID UNIVERSITY, PROBOLINGGO, INDONESIA
e-mail address: hamidelfath@gmail.com

MANABU OURA ²
 FACULTY of MATHEMATICS AND Physics, KANAZAWA UNIVERSITY
e-mail address: oura@se.kanazawa-u.ac.jp

(Received October 27, 2017)

(Accepted November 8, 2018)

TERWILLIGER ALGEBRAS OF SOME GROUP ASSOCIATION SCHEMES

ORIGINALITY REPORT

23%

SIMILARITY INDEX

23%

INTERNET SOURCES

8%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

www.laughingpanda.org

Internet Source

11%

2

ousar.lib.okayama-u.ac.jp

Internet Source

7%

3

oacs-dotlrn-conf2007.wu-wien.ac.at

Internet Source

1%

4

a100.gov.bc.ca

Internet Source

1%

5

www.dtic.mil

Internet Source

1%

6

mafiadoc.com

Internet Source

1%

7

doku.pub

Internet Source

1%

8

www.slideshare.net

Internet Source

<1%

9

idoc.pub

<1%

10

Martinez de Castilla, I.O.. "Cellular Binomial Ideals. Primary Decomposition of Binomial Ideals", Journal of Symbolic Computation, 200010

Publication

<1%

11

compgroups.net

Internet Source

<1%

12

Mark B. Yunker, Robie W. Macdonald, Lloyd R. Snowdon, Brian R. Fowler. "Alkane and PAH biomarkers as tracers of terrigenous organic carbon in Arctic Ocean sediments", Organic Geochemistry, 2011

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off