

ISSN 0257-8050

POLLUTION RESEARCH

EM INTERNATIONAL

VOL. 40 (2) : 2021

POLLUTION RESEARCH

VOL. 40 (2) : 2021

CONTENTS

- 1–8 **Stuđt Statistical Analysis of Air Pollution in Zarqa City - Jordan**
—*Sana'a Odat*
- 9–20 **Community-based Environment Management - A Pilot Study In Phu Dong Commune, Gia Lam District, Hanoi, Vietnam**
—*Pham Thi To Oanh and Tran Ngoc Thin*
- 21–32 **Environmental Biomonitoring of the Aquatic Bodies in Metro Manila, Philippines with Respect to the Genotoxicity Potential using *in vivo* mouse bone marrow micronuclei test and heavy metal analysis**
—*Mikaella Justin Umali and Zeba F Alam*
- 33–38 **An Overview on the Possible Challenges on Waste Disposal During Covid-19 Pandemic**
—*Nyathi S. and Olowoyo JO*
- 39–42 **Environmental Study to the Effect of Some Types of Pesticide on *Rhizobium sp.* bacteria**
—*May Hameed Mohammad AL-Dehamee*
- 43–48 **Removal Pb²⁺ of Well Water Using Purolite C-100 Resin and Adsorption Kinetic**
—*Esthi Kusdarini, Dian Yanuarita Purwaningsih and Agus Budianto*
- 49–55 **Quantitative Analysis of Plastic Polymeric Pollution at the Shore of the Shatt Al-Arab River in Southern Iraq**
—*Alaa K. Jasim*
- 56–65 **Mapping of CO₂ Emissions to Improve the Function of Green Open Space (Rth) in Kediri City**
—*Lisanudin Rahman, Mohammad Razif, Samsudin Affandi, Mochamad Junaidi Hidayat, Satria Fadil Persada and Prawira Fajarindra Belgiawan*
- 66–71 **Correlation of Water Discharge With Water Quality Parameters Using Sem in Sepanjang Bridge Monitoring Station and Gunungsari Dam of Surabaya River, Indonesia**
—*Jenius Setio Insanno, Mohammad Razif, Samsudin Affandi, Mochamad Junaidi Hidayat, Satria Fadil Persada, Prawira Fajarindra Belgiawan*
- 72–80 **Agricultural Wastes as a Carbon or Nitrogen Source for Production of Bacterial Cellulose. A Mini Review**
—*Khulood Fahad Saud Alabbosh, Nur Hazlin Hazrin Chong and Aisha Nawaf Al balawi*
- 81–86 **Characteristics of Polluters in the Working Environment of Pt X Palembang City**
—*Ferly Oktriyedi, Irfannuddin, Ngudiantoro, M. Hatta Dahlan and Nurhayati*
- 87–92 **Monitoring of environmental pollution for Pb, Cd and Cr elements using the feathers of three species of birds**
—*Asaad Sh. M. Alhesnawi*
- 93–97 **Estimation of Some Plant Nutritive and Heavy Metals in Euphrates River at the Cities of Ramadi and Khalidiah**
—*Shaimaa Fatih Ali and Bashar Tareq AL-Shandah*
- 98–104 **Prediction of Water Quality Status in Surabaya River With Dynamic System Method At Sepanjang Bridge Monitoring Station, Surabaya**
—*Ahmad Imaduddin, Mohammad Razif, Samsudin Affandi, Mochamad Junaidi Hidayat, Satria Fadil Persada and Prawira Fajarindra Belgiawan*
- 105–109 **Vermi-biofiltration for Removal of Chemical and Biological Pollutants from Municipal Wastewater For Agriculture – A Review**
—*Biswas, Amit*

- 110–117 Ecological Services of Arbuscular Mycorrhizal Fungi- A Reappraisal
—*Ameeta Sharma, Anuja Vajpayee, Neha Gheek Batra, Kalpana, Anjali Bhardwaj and P.C. Trivedi*
- 118–122 The Effect of Ajwa Dates Fruit Extract on Ovarian Follicle Development in Female Rats Exposed to Arsenic
—*Herdhika A.R. Kusumasari, Harwin H Desyanti, Mahanutabah H. Qurniatillah and Eviana Norahmawati*
- 123–126 Evaluating the Yield and Protein Content of Fodder Grass Under Paper Mill Effluent Irrigation and Solid Wastes Amendment
—*M. Suguna Devakumari (Assistant Professor)*
- 127–133 Physico-chemical Analysis of Contaminated Soil and Effluent Collected from Tannery Industries in Dindigul
—*Nirmala T., M.R. Delphine Rose, K. Jency Priya and B. Vijayabharathi*
- 134–141 Establishment of Water Quality Index in Different Waters of Theni and Dindigul Districts
—*T. Nirmala, Arockiam Thaddeus, Anita Kannagi, V. Santhi, S. Catherine P. Alexander, C. Sagaya Rani, M.R. Delphine Rose, V. Jemima Florence Borgia, S. Iruthaya Kalai Selvam, L. Merlin Dayana, A. Irudaya Lourdu Gladis, S. Devashanthi, P. Pushpa and R. Medona Mary*
- 142–145 Bio Removal of Arsenic Metals from Water Solution by *Bacillus Subtilis* sp.
—*Hazim A. Walli and Noor Ali Mohammed*
- 146–153 Physico Chemical Analysis of Coffee Effluent Before and After Treatment with *Tridax Procumbens* and its Impact on the Growth of Tomato Plant *Solanum Lycopersicum*
—*Nirmala T., K. Jency Priya and M.R. Delphine Rose*
- 154–157 Using Reduced Grapheme Oxide to Remove Arsenic Metals from the Environment
—*Hazim A. Walli and Jehan Adnan Abed*
- 158–162 Heavy Metals and its Pollution Leading to Resistance in Bacteria of Surface Waters: A Review
—*Richa Bhardwaj and Vigil Wilson*
- 163–169 Relationship of Water Quality with Phytoplankton Abundance in Kenjeran Coastal Waters, Surabaya, East Java, Indonesia
—*Anjani Arum Mahenda, Putu Angga Wiradana, Raden Joko Kuncoroningrat Susilo³, Trisnadi Agoes Soegianto, Arif Nur Muhammad Ansori and Novia Arinda Pradisty*
- 170–178 Screening Microorganisms for Phosphorus Removal in Saline Wastewater
—*Rafitah Hasanah, Tsuyoshi Imai, Ariyo Kanno, Takaya Higuchi, Masahiko Sekine and Koichi Yamamoto*
- 179–183 *Kocuria* species Identified as Lead Resistant Bacteria Isolated from Water Bodies of Udaipur Region
—*Garima Verma, Nadim Chishty and Sushma Jain*
- 184–191 Study of Different forms of Microbial Fuel Cell Technology (Mfc) for Generation of Electricity from Dairy Effluent Using A Microbial Consortium
—*Leena P. Pathak, V.B. Gaikwad and Sucheta Patil*
- 192–197 Geochemical Investigation of Hot Spring Water in Sikkim Himalaya: A Baseline Approach
—*Banashree Deori, Rakesh K. Ranjan and Shruti Dutta*
- 198–203 The Role of *Dendrobium* sp. as Lead Absorbent on the Surabaya Protocol Roads, Indonesia
—*Hamidah, Sucipto Hariyanto, Kartika K. Mutmainah and Manikya Pramudya*
- 204–206 Role of Soil Organic Matter in Plant Growth for Horizontal Flow Subsurface Constructed Wetlands for Waste Water Treatment
—*Arvind Kumar Swarnakar and Samir Bajpai*
- 207–216 Heavy Metal Tolerance Mechanisms in Microorganisms - A Review
—*Preeti Kanwar, Nikita Goswami and Tulika Mishra*

- 217–229 Occurrence of Antibiotic and Antibiotic Gene Resistant Contaminants in the Water Environment
—*Preeti Gahlot, Dimple and Rajesh Dhankhar*
- 230–236 The Utilization of Fruit Waste Extract in Pasar Induk Osowilangun Surabaya (PIOS) Indonesia as a Raw Material of Ethanol Bioconversion by *Saccharomyces Cerevisiae*
—*Nita Citrasari, Irma Ardiana Puspita and Agus Supriyanto*
- 237–239 Physicochemical and Bacterial Analysis of Textile Effluents from Kanchipuram, Tamilnadu
—*R. Kowsalya, K. Sujatha, Karthikeyan Sekar, Deepa Nagaraj and Sindhu Rajesh*
- 240–244 Determination of Lead and Cadmium in Edible Wedge Clam (*Donax Faba*) Collected from North and South Coasts of Sumenep, East Java, Indonesia
—*Qurrotul Aini Wasilah, Rizka Efi Mawli, Mayadita Dwi Sani, Agoes Soegianto, Putu Angga Wiradana and Novia Arinda Pradisty*
- 245–252 Microplastics: Pollutant of Emerging Concern- What Do We Know So Far? - A Review
—*Kshitij Upadhyay and Samir Bajpai*
- 253–258 Surveillance of *Vibrio* and *Blue-green Algae* in Intensive System of Pacific White Shrimp (*Litopenaeus vannamei*) in Situbondo Regency, East Java, Indonesia
—*Muhammad Inas Riandi, Raden Joko Kuncoroningrat Susilo, Mayadita Dwi Sani, Agnes Yuantin Maharani, Agoes Soegianto and Putu Angga Wiradana*
- 259–262 An Environmental Friendly Alternative to Chemical Surfactants by Exploring Biosurfactant Produced by *Acinetobacter baumannii* Isolated from Diesel Oilcontaminated Soil
—*Reshmi Gopalakrishnan, K. Madhanasundareswari and B. Thamarai Selvi*
- 263–267 Green Synthesis of Mild Steel Corrosion Inhibitor in a Salt Solution Made from Pepper Fruit (*Piper Nigrum Linn*), Areca Fruit (*Areca Cathecu Linn*) and Tea Leaves (*Cammellia Sinensis L. Kuntze*)
—*Siswanto and Djony Izak Rudyardjo*
- 268–277 Development of Inorganic Composites for Water Defluoridation
—*Praveen Tyagi and R.P. Singh*
- 278–283 Identification of Microplastic Content in Refilled Drinking Water in the District of Gunung Anyar Surabaya, Indonesia
—*Firra Rosariawari, Muhammad Nashruddin Abdulloh and Nurina Fitriani*
- 284–288 Potential of Tofu Waste as Biogas with the Addition of Manure and Sawdust
—*Nurina Fitriani, Firra Rosariawari and Andnurul Qomariyah*
- 289–295 Environmental Management Effort in Minimalist-styled Mosque With Audial Comfort and Noise Management
—*Mochamad Junaidi Hidayat, D.P.E. Laksmiyanti and Nurina Fitriani*
- 296–300 The combination of Bagasse and Chitosan as a Membrane in Elimination of Crossflow Batik Waste Pollutant Parameters
—*Firra Rosariawari, Fatia Hadiana and Wahid Dianbudiyanto*
- 301–306 Performance Evaluation of Electro Coagulation Process for Treatment of Groundwater
—*Munaf D. F. Al-Aseebee, Hameed R. Alamery and Ali S.J.*

Pollution Research Editorial Advisory Board

ISSN: 0257-8050

Advisors and Patrons

Prof. (Dr.) R.K.Trivedy, Pune, India

Email id: rktrivedy@gmail.com

Tel: 91-20-46745119

Mobile No: 9975703363

Former Professor and Head

Department of Environmental sciences

University of PUNE

C101-Prakriti

Balewadi, PUNE 411045,India

Chief Editor

1. Prof. Dr. Duangrat Inthorn

Deputy Dean for Research- Faculty of Public Health, Mahidol University

Mahidol University, Bangkok, Thailand

www.mahidol.ac.th

Contact: Tel/Fax: (662)354-8558

Email id: duangrat.int@mahidol.ac.th

2. Prof. Dr.V.Arutchelvan

Professor of Civil Engineering

Director, Centre for Skill Development

Annamalai University

Annamalainagar-608002

Tamilnadu, India

Mobile No: +91-9994773966

Email id- arul.au@gmail.com

EDITORIAL ADVISORY BOARD

1. Dr. Marck Bricka, Professor, Mississippi State University, U.S.A	bricka@che.msstate.edu
2. Dr. I.C. Onyema, Professor, Deptt. of Marine Sciences, University of Lagos, Akoka, Lagos, Nigeria	ionyema@unilag.edu.ng
3. Dr. Michael Green, Prof. Emeritus, Civil and Environment Engg. Technion, I.I.T., Haifa 32000, Israel	agmgreen@technion.ac.il
4. Dr. L.O. Chukwu, Professor, Department of Marine Sciences, University of Lagos, Nigeria	obinnalcunilag@yahoo.com
5. Dr. Prof. N.C. Gupta, Professor, Deptt. of Environmental Science and Technology, GGIP Univ., New Delhi	ncgupta@ipu.ac.in
6. Dr. S.I. Kolesnikov, Professor, Academy of Biology and Biotech. D.I. Ivanovsky, Southern Federal Univ. Russia	kolesnikov@sfedu.ru
7. Dr. Prof. J. Narayan, Professor and Head, Department of Environmental Science, Kuvempu University, Karnataka, India	narayanaj@kuvempu.ac.in
8. Dr. Neeka Jacob, Head, Research and Development at Petroleum Technology, PTDF, Abuja, Nigeria	neekajacob@uph.ng

9. Dr. A.H. Subratty, Prof. and Dean, University of Mauritius, Mauritius	deanfmhs@uom.ac.mu
10. Dr. K. Mathew, Director National Small Wind Turbine Test Centre, Murdoch University, Australia	kuruwilla@murdoch.edu.au
11. Dr. T. Bahorun, Professor, Department of Biosciences, Mauritius University, Mauritius	tbahorun (@) uom.ac.mu
12. Dr. C.K.Kale, Dean, Krishna Institute of Allied Sci., K.I.M.S., Deemed to be Univ., Karad, M.S., India	deankibb@kimskarad.in
13. Dr. Prof. Azni Idris, Deputy Vice Chancellor (Res. & Innovation), Universiti Putra Malaysia, Malaysia	azni@upm.edu.my
14. Dr. Prof. D.J. Lee, Professor, Department of Chemical Engineering, National Taiwan University, Taiwan	djlee@ntu.edu.tw
15. Dr. Margaret Greenway, Professor Griffith University, Australia	m.greenway@griffith.edu.au
16. Dr. Prof. Ir. Diana Arfiati,, Professor, School of Fisheries and marine Sciences, University of Brawijaya, Indonesia	d-arfiati@ub.ac.id
17. Dr. A. Giacometti, Professor, Department of Environmental Science, University of Venice, Venice, Italy	giacomet@unive.it
18. Dr. Prof. A.K. Dikshit, Professor, Centre for Environmental Science, I.I.T., Mumbai, India	dikshit@iitb.ac.in
19. Dr. Prof. Ajit Pratap Singh, Professor and Dean, Birla Institute of Technology, Pilani, Rajasthan, India	aps@pilani.bits-pilani.ac.in
20. Dr. Prof. S.A. Abbasi, Professor, Deptt. of Environmental Science, Pondicherry University, Puducherry, India	abbasi.cpee@gmail.com
21. Dr. Prof. A.R. Ghosh, Professor, Department of Environmental Science, Burdwan University, Burdwan, India	apurbaghosh2010@gmail.com
22. Dr. Prof. Deenbandhu Sahoo, Professor, Department of Botany, University of Delhi, India	dbsahoo@hotmail.com
23. Dr. Prof. D.P. Singh, Professor, Dept. of Environmental Sciences, Dr. B.B.A. University, Lucknow, India	dpsingh_lko@yahoo.com
24. Dr. Prof. Toan Vu, Duc, Professor, Thuyli University, Hanoi, Vietnam	toanvd@wru.vn
25. Dr. Suresha, Huta, Senior Environmental Consultant, Environmental Company, Saudi Arabia	vaishnavisureshg@yahoo.com
26. Dr. Prof. T.N. Singh, Professor (Presently Vice-Chancellor, Kashi Vidyapeeth Univ., Varanasi) Department of Geology, I.I.T, Mumbai, India	vc@mgkvp.ac.in
27. Dr. Vijaya Ilango, Associate Professor Department of Chemistry, Birla Institute of Technology and Science, UAE	vilango@dubai.bits-pilani.ac.in
28. Dr. Prof. R.M. Naraynan,, Professor and head Dept. of Civil Engineering, Dr. M.G.R. Deemed to be Univ., Chennai, India	narayanan.rm@drmgrdu.ac.in
29. Dr. Kudrat-E-Khuda (Babu), Assoc. Prof. & Head, Dept. of Law, Daffodil Intern. Univ., Bangladesh	kekbabu@gmail.com
30. Dr. V.N. Janakirajan, Assoc. Prof., Department of Biochemistry, Govt. Medical College, Erode, India	vnjngeetha@gmail.com
31. Dr. Suresha Gududappa, Senior Environmental Scientist, Saudi Arabia	suresh@ebe.com.sa

THE EFFECT OF AJWA DATES FRUIT EXTRACT ON OVARIAN FOLLICLE DEVELOPMENT IN FEMALE RATS EXPOSED TO ARSENIC

HERDHIKA A.R. KUSUMASARI^{1*}, HARWIN H DESYANTI^{1,2}, MAHANUTABAH H. QURNIATILAH^{1,3} AND EVIANA NORAHMAWATI⁴

¹Master of Midwifery Program, Brawijaya University, Malang, Indonesia

²Midwifery Program, Faculty of Health, Nurul Jadid University, Probolinggo, Indonesia

³Harapan Kita Women and Children Hospital, Jakarta, Indonesia

⁴Department of Anatomic Pathology, Faculty of Medicine, Brawijaya University, Malang, Indonesia

ABSTRACT

Arsenic trioxide (As_2O_3) is a form of inorganic arsenic which is very toxic in the environment. Arsenic exposure can cause reproductive health problems, one of which is the ovaries, by interfering with the follicle development. Arsenic exposure causes oxidative stress in the ovaries due to the accumulation of excessive *Reactive Oxygen Species* (ROS) and an imbalance between ROS and antioxidants in the body. The powerful antioxidants contained in dates can protect the effects of free radicals due to arsenic exposure. The group in this study was divided into a negative control group (without arsenic exposure), positive control (3 mg / kg BW arsenic), T1 (arsenic 3 mg / kg BW + date extract 2mL / kg BW), T2 (arsenic 3 mg / kg BW + date extract 4ml / kg BW), and T3 (arsenic 3 mg / kg BW + date extract 8ml / kg body weight). The administration of Ajwa date extract could significantly provide antioxidant protection in the treatment group with an increase in follicle count on this study.

KEY WORD : Arsenic trioxide, Arsen, Ovary, Free Radical, *Reactive Oxygen Species*, Oxidative Stress

INTRODUCTION

Exposure to heavy metals, one of which is arsenic, which is a group *metalloid*, can cause reproductive dysfunction in women (Mendola *et al.*, 2008), even residual heavy metal exposure in the body can affect women's reproductive health (Bloom *et al.*, 2011). Arsenic toxicity has become a major concern due to increased contamination of exposure to this compound in the air, water, and soil (Flora, 2014). *The Agency for Toxic Substances and Disease Registry* United States (ATSDR) has designated arsenic as No. 1 on the Priority List of Hazardous Substances over the lead, mercury and *Vinyl Chloride* (ATSDR, 2019) (Shen *et al.*, 2013). Currently, the number of humans exposed to arsenic through the air, food, water, and soil is increasing all the time. Arsenic

toxic effects are very common in developing countries. Studies on the toxicity of arsenic exposure in women have been carried out in several countries. In a study conducted on women in Taiwan by Lei *et al.* (2015) arsenic levels in infertile women were higher than in pregnant women. A case study conducted on women in China who were exposed to arsenic found that arsenic can cause ovarian insufficiency / *Premature Ovarian Insufficiency* (Pan *et al.*, 2020). A drastic decrease in the number of follicles due to environmental factors such as heavy metal exposure, namely arsenic, can result in early menopause (Kumar *et al.*, 2012).

The effect of arsenic exposure on ovarian organ can cause damage to ovarian morphology while folliculogenesis affects follicular regression. Arsenic exposure causes a decrease in the number of follicles

including primordial, primary, secondary and *de Graaf* follicles and increases the number of atretic follicles (Chattopadhyay and Ghosh, 2010; Akram *et al.*, 2010; Mehta and Hundal, 2016; Khatun *et al.*, 2018; Yu *et al.*, 2019).

Many studies have examined the benefits of dates fruit for female's reproductive health because they are believed to have an effect on oocyte quality, sperm and ovum interactions, implantation and early embryonic development through antioxidant mechanisms. (Abdi *et al.*, 2017; Saryono *et al.*, 2018). The antioxidant properties of Ajwa dates are expected to be able to suppress free radicals through *scavenger* pathway, which have an effect to reduce disease proliferation due to the strong antioxidant content in Ajwa dates (Ahmed *et al.*, 2016; Al-Yahya *et al.*, 2016). Therefore, in this study, we are interested in conducting studies on the effects of Ajwadata extract (*Phoenix dactylifera* L.) on the female reproductive system, especially on follicle development in female Wistar rats (*Rattus norvegicus*) and the lack of research on the effects of Ajwa date palm extract in preventing exposure arsenic metal toxicity.

MATERIALS AND METHODS

Chemical

Arsenic powder (As_2O_3) produced by LobaChemie as much as 500 mg was dissolved in 1000 mL *Normal Saline* 0.9% then stirred with a *magnetic stirrer* on heating at 50 °C for 3-4 hours to make the arsenic powder completely dissolve.

Preparation of Extract The

Types of dates used for the study were 400 grams of Ajwa dates. Selection of dates by selecting dates that are old, not rotten, separated from the seeds, then thinly sliced and the flesh of the dates is taken. The extract making refers to previous research, namely research by Vayalil (2002) and Al-Rasheed *et al.*, (2015) with slight modifications. The flesh of the ajwa dates that have been separated from the seeds, mashed with a *mortar* and *pestle*, then added 1200 ml of water (ratio 1: 3, g/ml). The thick date palm extract was centrifuged at 4°C for 15 minutes at a speed of 10,000g, then the supernatant was taken.

Animals and Ethics Statement

Research procedures have been accepted by the Health Research Ethics Commission, Faculty of Medicine, University of Brawijaya. In this study,

there were 5 treatment groups where there were 5 repetitions for each group. Negative control was rats given *Normal Saline* and not given date palm extract and arsenic, positive control rats exposed to arsenic 3mg/kg BW/day, treatment 1, 2 and 3 rats given date extracts with 3 different doses (2ml / KgBW / day; 4ml/KgBB/day; 8ml/KgBB/day), then exposed to 3 mg/KgBB / day arsenic for 30 days. The research was conducted at the Laboratory of Bioscience, Anatomical Pathology and Biochemistry Laboratory, Faculty of Medicine, University of Brawijaya.

Vaginal Smears Collection

After 30 days of treatment, a swab was performed to determine the rat oestrous cycle. Mice in the proestrus phase were terminated. The swab samples were stained with *methylene blue* and observed the morphology of the epithelial cells under a microscope (Sezer *et al.*, 2020).

Histological Examination

Number of Follicles. Ovarian samples were stained with *Hematoxylin Eosin (H&E)*, criteria for follicular classification based on Myers *et al.* (2004). The results of the staining of the sample were then viewed under a microscope *Dot slide Olympus XC 10* and the number of follicles was calculated using *Dot Slide software*.

Data Analysis

Data were analyzed using software *SPSS 25.0*

RESULTS AND DISCUSSION

Note: The mean number of ovarian follicles based on the treatment group. On the mean \pm SD, if it contains different letters, it means that there is a significant difference ($p \leq 0.05$) and if it contains the same letter, it means that there is no significant difference ($p \geq 0.05$).

The number of primordial follicles showed a significant result with p -value = 0.043. There was a decrease in the number of primordial follicles in the arsenic-exposed only group (2.4 ± 2.607^a) compared to the healthy or negative control groups (2.6 ± 1.949^a), but the decrease was not statistically significant. In the group of mice given Ajwa date extract at a dose of 4 ml/kg BW, there was a significant increase in the number of primordial follicles compared to the positive control group that was given arsenic alone.

The same thing happened to the number of primary follicles, all treatment groups showed significant results statistically with a value of $p = 0.029$. Although in the positive control group (2.4 ± 2.607^a) there was a decrease in the number of secondary follicles, the decrease was not significant compared to the negative control group ($2.6 \pm 1,949^a$) statistically. Ajwa date extract caused a protective effect on the number of primary follicles in female rats exposed to arsenic with the highest dose of date extract, 8 ml/kg BW. In treatment group 3 ($3.0 \pm 1,000^{ab}$) there was a significant increase in the number of primary follicles compared to the arsenic exposed group (1.6 ± 0.547^a).

In the number of secondary follicles, the group of mice exposed to arsenic (1.2 ± 1.095^a) had a decrease in number but not significantly compared to the group of rats that were either healthy or not exposed to arsenic (2.4 ± 1.140^a). In the treatment group given date palm extract 8 ml/kg BW or Treatment 3 ($2.0 \pm 0,707^a$) there was an increase in the number of secondary follicles compared to the positive control group (1.2 ± 1.095^a), the mean value of the P3 group approached the negative control group (2.4 ± 1.140^a). However, there was an increase in the number of secondary follicles with a mean value that exceeded the group of healthy rats that occurred in the treatment group given date extract 4 ml/kg BW ($4.2 \pm 3,768^a$).

Then the number of early antral follicles also decreased the number of early antral follicles in the positive control group (0.8 ± 0.836^a) compared to the negative control group/only given Normal saline (1.2 ± 0.447^a), but the decrease was not significant. There were interesting results in the group of mice given date palm extract at a dose of 4 ml/kg BW (1.2 ± 0.446^a) and 8 ml/kg BW (1.2 ± 0.447^a) where the value of the number of early antral follicles increased, and the mean value was the same as healthy mice group (1.2 ± 0.447^a), whereas, at the smallest dose of date extract, 2 ml/kg BW, there was

an increase in the number of early antral follicles (2.0 ± 1.000^a) compared to the mice exposed to arsenic alone.

In addition to the number of follicles mentioned in this study, researchers also counted the number of antral or *Graafian follicles* that was published. The results obtained were that there was a significant decrease in the number of *Graafian follicles* in mice exposed to arsenic and date palm extracts can increase the number of *Graafian follicles* in female rats exposed to arsenic (Kusumasari *et al.*, 2020).

The decrease in the number of follicles in the group of rats exposed to arsenic in this study was the same as previous research, namely Mehta and Hundal (2016) that rats exposed to arsenic caused problems in folliculogenesis. The decrease in the number of follicles occurs due to the mechanism of arsenic toxicity in various ways so that it affects the disruption of folliculogenesis. One of them is arsenic exposure causes an increase in *Reactive Oxygen Species (ROS)* which results in oxidative stress on the ovaries (Flora, 2011; Sun *et al.*, 2016). The increase in ROS is usually followed by an increase in lipid peroxidase and MDA which are biomarkers of oxidative stress. Homeostatic changes in the body, such as increased levels of ROS due to arsenic exposure, also alter the activity of antioxidant enzymes, which act as a first-line defence against free radicals. Inhibition of antioxidant enzymes will cause excessive levels of superoxide anions (Banerjee *et al.*, 2014), excessive accumulation of superoxide anions can cause follicular regression (Khatun *et al.*, 2018). Arsenic exposure also inhibits the peroxidase enzyme in the follicular fluid so that it interferes with folliculogenesis due to high levels of free radicals (Chattopadhyay and Ghosh, 2010). Apart from going through the ROS pathway, arsenic toxicity can damage DNA and damage ovarian cells (Akram *et al.*, 2009). A drastic decrease in the number of follicles caused by environmental factors such as heavy metal exposure, namely arsenic, can

Table 1. Effect of Ajwa Dates Extract on Ovarian Follicles Count of Female Rats.

Follicle	Negative Control Mean±SD	Positive Control (arsenic) Mean±SD	Treatment 1 (arsenic+date extract 2 ml/kgBB) Mean±SD	Treatment 2 (arsenic+date 4 ml/kgBB) Mean±SD	Treatment 3 (arsenic+date 8 ml/kgBB) Mean±SD
Primordial	2.6±1.949 ^a	2.4±2.607 ^a	3.0 ±2.645 ^a	7.0±3.082 ^{ab}	13.2±12.336 ^a
Primer	2.6±1.516 ^a	1.6±0.547 ^a	1.2 ±0.447 ^a	1.8±0.447 ^a	3.0±1.000 ^{ab}
Sekunder	2.4±1.140 ^a	1.2±1.095 ^a	1.2 ±0.836 ^a	4.2±3.768 ^a	2.0±0.707 ^a
Early Antral	1.2±0.447 ^a	0.8±0.836 ^a	2.0 ±1.000 ^a	1.2±0.446 ^a	1.2±0.447 ^a

result in premature menopause in women (Kumar *et al.*, 2012).

The effect of arsenic on the endocrine system is also one of the causes of disturbances in follicular development. Arsenic exposure indirectly interferes with the regulation of the ovarian pituitary axis so that it affects decreasing levels of the *Follicle Stimulating Hormone* (FSH) hormone and decreasing levels of the steroid estrogen hormone where the action of this hormone is needed in the folliculogenesis process. If there is inhibition of hormones that affect follicle development, it can result in low follicle regression and ovarian mass (Khatun *et al.*, 2018). Therefore, in this study, the group of mice exposed to arsenic alone had a decreased number of follicles compared to the healthy group of mice.

The preventive effect of Ajwa date fruit extract in this study is due to the antioxidant content contained in dates, namely melatonin, carotene, phenolic compounds, flavonoids and contains vitamins C, E and vitamin A. Phenolic compounds contained in dates are routine, catechin, *caffeic acid*, *gallic acid*, *quercetin*, *luteolin* and *apigenin* (Chaira *et al.*, 2007; Al-Farsi and Lee, 2008; Ben Abdallah *et al.*, 2009; Ali *et al.*, 2015; Eid *et al.*, 2015; Khalid *et al.*, 2017). Apart from phenolic compounds, a study conducted by Chao (2007) proved that dates contain estradiol.

The work of antioxidants in Ajwa date extract by inhibiting lipid peroxidase so that it can act as an antioxidant that stops reactions associated with oxidative stress (Zhang *et al.*, 2013). The antioxidants contained in Ajwa dates can stop the bonding reactions associated with oxidative stress by reducing the formation of ROS (Ben Abdallah *et al.*, 2009). The phenolic content of Ajwa dates, with their redox properties, can act as a *scavenger* ROS, neutralize free radicals, and break down peroxides (Kchaou *et al.*, 2014). So that in this study proved that Ajwa date extract had a protective effect on ovarian follicles in female rats exposed to arsenic.

CONCLUSIONS

In this study, it can be concluded that arsenic exposure can affect the number of ovarian follicles and Ajwa date extract which contains high antioxidants can protect ovarian follicles from damage caused by arsenic exposure.

ACKNOWLEDGEMENT

Thanks to the laboratory analysts of Bioscience, Pathology Anatomy, Biochemistry and Laboratory of the Center for Biological Sciences (LSIH), Brawijaya University.

REFERENCES

- Ben Abdallah, F. 2009. Effects of date seed oil on testicular antioxidant enzymes and epididymal sperm characteristics in male mice. *Andrologia*. 41(4) : 229-234.
- Abdi, F., Roozbeh, N. and Mortazavian, A.M. 2017. Effects of date palm pollen on fertility: Research proposal for a systematic review. *BMC Research Notes*. 10(1) : 1-5.
- Ahmed, A. et al., 2016. Nutritional probing and HPLC profiling of roasted date pit powder. *Pakistan Journal of Nutrition*. 15(3) : 229-237.
- Akram, Z. 2010. Adverse effects of arsenic exposure on uterine function and structure in female rat. *Experimental and Toxicologic Pathology*. 62(4): 451-459. Available at: <http://dx.doi.org/10.1016/j.etp.2009.07.008>.
- Akram, Z. 2009. Genotoxicity of sodium arsenite and DNA fragmentation in ovarian cells of rat. *Toxicology Letters*. 190(1) : 81-85.
- Al-Farsi, M.A. and Lee, C.Y. 2008. Optimization of phenolics and dietary fibre extraction from date seeds. *Food Chemistry*. 108(3) : 977-985.
- Al-rasheed, Nouf M. 2015. Aqueous Date Flesh or Pits Extract Attenuates Liver Fibrosis via Suppression of Hepatic Stellate Cell Activation and Reduction of Inflammatory Cytokines, Transforming Growth Factor- β 1 and Angiogenic Markers in Carbon Tetrachloride-Intoxicated Rats. *Hindawi Publishing Corporation Evidence-Based Complementary and Alternative Medicine*, Volume 201, p.19 pages.
- Al-Yahya, M. 2016. 'Ajwa' dates (*Phoenix dactylifera* L.) extract ameliorates isoproterenol-induced cardiomyopathy through downregulation of oxidative, inflammatory and apoptotic molecules in rodent model. *Phytomedicine*. 23(11) : 1240-1248.
- Ali, E., Sulieman, A.M.E., Elhafise, I.A.A. and Abdelrahim, A.M. 2015. Comparative Study on Five Sudanese Date (*Phoenix dactylifera* L.) Fruit Cultivars Comparative Study on Five Sudanese Date (*Phoenix dactylifera* L.) Fruit Cultivars. , 2012(NOVEMBER), pp.1245-1251.
- Banerjee, Sayani, 2014. Female Reproductive Aging Is Master-Planned at the Level of Ovary Franks, S., (ed.). *PLoS ONE*. 9(5) : e96210. Available at: <https://dx.plos.org/10.1371/journal.pone.0096210> [Accessed: 16 January 2020].
- Bloom, M.S. 2011. Associations between blood metals

- and fecundity among women residing in New York State. *Reproductive Toxicology*. 31(2) : 158-163. Available at: <http://dx.doi.org/10.1016/j.reprotox.2010.09.013>.
- Chaira, N., Ferchichi, A., Mrabet, A. and Sghairoun, M., 2007. Chemical Composition of the Flesg and the Pit of Date Palme Friut 2007.Pdf. *Pakistan Journal of Biological Science*. 10(13) : 2202-2207.
- Chao, C.T. and Krueger, R.R., 2007. The date palm (*Phoenix dactylifera* L.): overview of biology, uses, and cultivation. *Hort Science*. 42(5) : 1077-1082.
- Chattopadhyay, S. and Ghosh, D., 2010. Role of dietary GSH in the amelioration of sodium arsenite-induced ovarian and uterine disorders. *Reproductive Toxicology*. 30(3) : 481-488. Available at: <http://dx.doi.org/10.1016/j.reprotox.2010.05.002>.
- Chuan-Rui Zhang, Saleh A. Aldosari, Polana S. P. V. Vidyasagar, Karun, M. Nair, and M.G.N. 2013. Antioxidant and Anti-inflammatory Assays Con firm Bioactive Compounds in Ajwa Date Fruit. *Journal of Agricultural and Food Chemistry*.
- Eid, N. 2015. Impact of palm date consumption on microbiota growth and large intestinal health: A randomised, controlled, cross-over, human intervention study. *British Journal of Nutrition*, 114(8) : 1226-1236.
- Flora, S.J.S., 2011. Arsenic-induced oxidative stress and its reversibility. *Free Radical Biology and Medicine*, 51(2) : 257-281. Available at: <http://dx.doi.org/10.1016/j.freeradbiomed.2011.04.008>.
- Flora, S.J.S., 2014. *Handbook of arsenic toxicology*, Academic Press.
- Kchaou, W., Abbès, F., Attia, H. and Besbes, S., 2014. In vitro antioxidant activities of three selected dates from Tunisia (*Phoenix dactylifera* L.). *Journal of Chemistry*, 2014.
- Khalid, S. 2017. A review on chemistry and pharmacology of Ajwa date fruit and pit. *Trends in Food Science and Technology*. 63(September), pp.60-69. Available at: <http://dx.doi.org/10.1016/j.tifs.2017.02.009>.
- Khatun, S. 2018. *Spirulina platensis* ameliorates arsenic-mediated uterine damage and ovarian steroidogenic disorder. *Facets*. 3 (1) : 736-753.
- Kumar, M. 2012. Chromosomal abnormalities and oxidative stress in women with premature ovarian failure (POF). *Indian Journal of Medical Research*. 135 (1) : 92-97.
- Kusumasari, H.A.R., Desyanti, H.H. and Mahanutabah, H. 2020. The Effect of Ajwa Dates Fruit Extract on Follicle Stimulating Hormone (FSH), Graafian Follicle and Endometrial Thickness in Female Rats Exposed to Arsenic. *Journal of Global Pharma Technology*. 12(08) : 122-130.
- Lei, H.L. 2015. Relationship between risk factors for infertility in women and lead, cadmium, and arsenic blood levels: A cross-sectional study from Taiwan. *BMC Public Health*. 15(1) : 1-11. Available at: <http://dx.doi.org/10.1186/s12889-015-2564-x>.
- Mehta, M. and Hundal, S.S. 2016. Effect of sodium arsenite on reproductive organs of female Wistar rats. *Archives of Environmental and Occupational Health*. 71 (1) : 16-25.
- Mendola, P., Messer, L.C. and Rappazzo, K. 2008. Science linking environmental contaminant exposures with fertility and reproductive health impacts in the adult female. *Fertility and Sterility*, 89(2 SUPPL.) : 81-94.
- Myers, M. 2004. Methods for quantifying follicular numbers within the mouse ovary. *Reproduction*. 127(5) : 569-580.
- Pan, W. 2020. A case-control study of arsenic exposure with the risk of primary ovarian insufficiency in women. *Environmental Science and Pollution Research*. 27(20) : 25220-25229.
- Saryono, S., Dwi, M. and Rahmawati, E., 2018. Effects of Dates Fruit (*Phoenix dactylifera* L.) in the Female Reproductive Process. 03(October), pp.1630-1633. Available at: https://www.academia.edu/33897003/EFFECTS_OF_DATES_FRUIT_PHOENIX_DACTYLIFERA_L_IN_THE_FEMALE_REPRODUCTIVE_PROCESS.
- Sezer, Z. 2020. Effects of vitamin E on nicotine-induced lipid peroxidation in rat granulosa cells: Folliculogenesis. *Reproductive Biology*. 20(1) : pp.63-74.
- Shen, S. 2013. Arsenic binding to proteins. *Chemical Reviews*. 113(10) : 7769-7792.
- Sun, H.J. 2016. Mechanisms of arsenic disruption on gonadal, adrenal and thyroid endocrine systems in humans: A review. *Environment International*. 95 : 61-68. Available at: <http://dx.doi.org/10.1016/j.envint.2016.07.020>.
- Vayalil, P.K. 2002. Antioxidant and antimutagenic properties of aqueous extract of date fruit (*Phoenix dactylifera* L. *Arecaceae*). *Journal of Agricultural and Food Chemistry*. 50(3) : 610-617.
- Yu, H. 2019. Sodium Arsenite Injection Induces Ovarian Oxidative Stress and Affects Steroidogenesis in Rats. *Biological Trace Element Research*. 189 (1) : 186-193.
-